

8-2001

On Preserving Structured Matrices Using Double Bracket Operators: Tridiagonal and Toeplitz Matrices

Kenneth Driessel
University of Wyoming

Irvin R. Hentzel
Iowa State University, hentzel@iastate.edu

Wasin So
San Jose State University

Follow this and additional works at: http://lib.dr.iastate.edu/math_pubs

 Part of the [Algebra Commons](#)

The complete bibliographic information for this item can be found at http://lib.dr.iastate.edu/math_pubs/156. For information on how to cite this item, please visit <http://lib.dr.iastate.edu/howtocite.html>.

This Article is brought to you for free and open access by the Mathematics at Iowa State University Digital Repository. It has been accepted for inclusion in Mathematics Publications by an authorized administrator of Iowa State University Digital Repository. For more information, please contact digirep@iastate.edu.

On Preserving Structured Matrices Using Double Bracket Operators: Tridiagonal and Toeplitz Matrices

Abstract

In the algebra of square matrices over the complex numbers, denotes Two problems are solved: (1) Find all Hermitian matrices M which have the following property: For every Hermitian matrix A , if A is tridiagonal, then so is (2) Find all Hermitian matrices M which have the following property: For every Hermitian matrix A , if A is Toeplitz, then so is

Keywords

structured matrices, tridiagonal, Toeplitz

Disciplines

Algebra | Mathematics

Comments

This article is published as Driessel, Kenneth R., Irvin R. Hentzel, and Wasin So. "On Preserving structured matrices using double bracket operators: Tridiagonal and Toeplitz matrices," JP Journal of Algebra, Number Theory and Applications, 1, no. 2, (2001): 87-114. Posted with permission.

ON PRESERVING STRUCTURED MATRICES USING DOUBLE BRACKET OPERATORS: TRIDIAGONAL AND TOEPLITZ MATRICES

KENNETH R. DRIESSEL

*Department of Mathematics, University of Wyoming,
Laramie, WY 82071-3036, U. S. A.*

IRVIN R. HENTZEL

*Mathematics Department, Iowa State University,
Ames, Iowa, 50011-2066, U. S. A.*

WASIN SO

*Department of Mathematics and Computer Science,
San Jose State University, San Jose, CA 95192-0103, U. S. A.*

Abstract

In the algebra of square matrices over the complex numbers, $[X, Y]$ denotes $XY - YX$. Two problems are solved: (1) Find all Hermitian matrices M which have the following property: For every Hermitian matrix A , if A is tridiagonal, then so is $[A, [A, M]]$. (2) Find all Hermitian matrices M which have the following property: For every Hermitian matrix A , if A is Toeplitz, then so is $[A, [A, M]]$.

1. Introduction

In this paper all matrices are square and complex and the set of all

2000 Mathematics Subject Classification: 15A04, 65F15.

Key words and phrases: structured matrices, tridiagonal, Toeplitz.

Communicated by Zhongshan Li

Received September 9, 2001

© 2001 Pushpa Publishing House

$n \times n$ complex matrices is denoted by $C^{n \times n}$. We use the Lie bracket notation, $[X, Y] = XY - YX$. We call the map $X \rightarrow [X, [X, M]]$ on $C^{n \times n}$ the *double bracket* map determined by M . Note that if A and M are Hermitian, then $[A, M]$ is skew-Hermitian and $[A, [A, M]]$ is Hermitian. In other words, if M is Hermitian, then the associated double bracket map is "Hermitian preserving." The following two problems are considered:

Problem A. Find all Hermitian matrices M with the "Jacobi preserving" property: For every Hermitian matrix A , if A is tridiagonal, then so is $[A, [A, M]]$.

Problem B. Find all Hermitian matrices M with the "Toeplitz preserving" property: For every Hermitian matrix A , if A is Toeplitz, then so is $[A, [A, M]]$.

We shall present the solutions, using elementary techniques only, in Sections 2 and 3 respectively.

Before presenting our solutions we want to explain why we chose to study these problems. The need to calculate the eigenvalues of large structured matrices leads to them. Consider the initial value problem with the following "double bracket" ordinary differential equation:

$$X' = [X, [X, M]], \quad X(0) = A,$$

where M and A are Hermitian matrices. The solution $X(t)$ of this problem has very interesting properties summarized in Proposition 1.2. These properties are known in literature. Chu and Driessel [9] considered the optimization problem: Given square Hermitian matrices A and M , find the matrix which is closest to M subject to the constraint that it is unitarily similar to A . They derived the (gradient) double bracket differential equation from this optimization problem. Bloch, Brockett and Ratiu [2] studied this differential equation and showed that if M is the diagonal matrix with diagonal entries $1, 2, \dots, n$, then the double bracket differential equation is closely related to the Toda flow and the QR algorithm. See Nanda [20], Brockett [3, 4, 5], Bloch [1], and Chu [11]. The

connection between the Toda flow and the QR algorithm is due to Symes [22, 23]. Deift, Nanda and Tomei [12] emphasized the use of isospectral flows to find eigenvalues. Chu [6] and Watkins [27] wrote survey articles concerning these matters. For description of the QR algorithm and its origins see, for example, Parlett [21] or Golub and Van Loan [19]. There are differential equations related to the Jacobi methods for finding eigenvalues; see Driessel [14, 15]. For descriptions of traditional Jacobi methods, see, for example, Parlett [21], Golub and Van Loan [19] or Wilf [28]. For more on the differential geometry of isospectral surfaces, see Tomei [25] and Driessel [17]. There are also differential equations for finding singular values; see Chu [7], Driessel [16], Deift, Demmel, Li and Tomei [13].

The following proposition shows that if a smooth vector field preserves a linear structure, then so does the corresponding flow. This result is an instance of more general results concerning vector fields tangent to manifolds. See, for example, Thorpe [24], Chapter 5 "Vector fields on surfaces; orientation."

Proposition 1.1. *Let V be a normed linear space over the real numbers and let U be a subspace of V . Let $F : V \rightarrow V$ be a continuously differentiable function satisfying $F(U) \subseteq U$. Let a be an element of U and let $x(t)$ be the solution of the initial value problem: $x' = F(x)$, $x(0) = a$. Then the solution curve $x(t)$ lies in U .*

Proof. Let W be a subspace of V which is a complement of U ; in symbols, $V = U \oplus W$. Let P_W and P_U be the linear projections of V onto W and U determined by this direct sum decomposition. Consider the initial value problem: $y' = F(P_U(y))$, $y(0) = a$. Let $y(t)$ denote the solution of this initial value problem. Note that $[P_W(y)]' = P_W(y') = P_W(F(P_U(y))) = 0$. Hence $P_W(y(t))$ is a constant. Since $P_W(y(0)) = P_W(a) = 0$, this constant is 0. Thus $y(t)$ is in U . It follows that $y(t)$ is a solution of the initial value problem that determines $x(t)$. By the uniqueness of the solution of this latter problem, we have $x(t) = y(t)$ and $x(t)$ is in U .

The next proposition shows that we can use the double bracket differential equation to find the eigenvalues of Hermitian matrices.

Proposition 1.2. *Let A and M be Hermitian matrices and let $X(t)$ be the solution of the following initial value problem (*):*

$$X' = [X, [X, M]], \quad X(0) = A.$$

Then this solution has the following properties:

(1) *For all t , $X(t)$ is a Hermitian matrix with the same eigenvalues as A .*

(2) *The solution $X(t)$ converges to a Hermitian matrix which commutes with M .*

(3) *If M is a diagonal matrix with distinct diagonal entries, then the solution $X(t)$ converges to a diagonal matrix with diagonal entries equal to the eigenvalues of A .*

Proof. It follows from Proposition 1.1 that the solution $X(t)$ is Hermitian for all t since double bracket map is Hermitian preserving. Define $K(t) = [X(t), M]$. Note $K(t)$ is skew Hermitian since $X(t)$ and M are Hermitian. We consider the following initial value problem (**):

$$Q' = QK, \quad Q(0) = I.$$

Claim. The solution $Q(t)$ of (**) is unitary for all t .

Note that $(QQ^*)' = Q'Q^* + Q(Q')^* = QKQ^* - QKQ^* = 0$ since K is skew Hermitian. Hence $Q(t)Q^*(t)$ is constant. This constant is I since $Q(0)Q(0)^* = I$.

Claim. The solution $X(t)$ of (*) satisfies $X(t) = Q(t)^* A Q(t)$ for all t .

Define $Y(t) := Q(t)^* A Q(t)$. Then

$$Y' = (Q^* A Q)' = (Q')^* A Q + Q^* A Q' = -KQ^* A Q + Q^* A Q K = [Y, K].$$

Thus, Y is a solution of the following initial value problem:

$$Y' = [Y, K], \quad Y(0) = A.$$

But X is also a solution of this initial value problem. Hence by uniqueness of such solutions, $X = Y$. Hence (1) is proved.

We shall regard $C^{n \times n}$ as a real vector space (with dimension $2n^2$) with the real inner product:

$$(P, Q) := \operatorname{Re}(\operatorname{Trace}(PQ^*)),$$

where $\operatorname{Re} z$ denotes the real part of the complex number z and $\operatorname{Trace}(R)$ denotes the trace of the matrix R . In terms of coordinates $(P, Q) = \sum_{i,j} (\operatorname{Re}(P_{ij}) \operatorname{Re}(Q_{ij}) + \operatorname{Im}(P_{ij}) \operatorname{Im}(Q_{ij}))$, where $\operatorname{Im} z$ denotes the imaginary part of a complex number z . We have

$$\frac{d}{dt} \left(\frac{1}{2} \|X(t) - M\|^2 \right) = -([X, M], [X, M]) \leq 0 \quad (\dagger)$$

since

$$\begin{aligned} \frac{d}{dt} \frac{1}{2} (X - M, X - M) &= (X - M, X') \\ &= (X - M, [X, [X, M]]) \\ &= ([X, X - M], [X, M]) \\ &= -([X, M], [X, M]). \end{aligned}$$

We use the fact that $(PQ, R) = (Q, P^*R) = (P, RQ^*)$ and $([P, Q], R) = (Q, [P^*, R])$. We see that the solution $X(t)$ remains bounded since $\|X(t) - M\| \leq \|A - M\|$. We also see from (\dagger) that $X(t)$ converges to a matrix Y which satisfies $[M, Y] = 0$. Thus, $X(t)$ satisfies property (2).

Now assume that $M = \operatorname{Diag}(d_1, d_2, \dots, d_n)$ is a diagonal matrix with distinct diagonal entries. Then $[M, Y]_{ij} = (d_i - d_j)Y_{ij}$. Hence $[M, Y] = 0$

implies that Y is a diagonal matrix. Thus we see that property (3) follows from properties (1) and (2).

There are computer softwares available for numerically solving differential equations $X' = F(X)$. The evaluations of the function F usually constitute most of the numerical work. For the double bracket differential equation we have $F(X) = [X, [X, M]]$. Generally the evaluation of the double bracket requires $O(n^3)$ operations. We can reduce this work if the double bracket operation preserves structure. For example, if X are tridiagonal, then the bracket computation only requires $O(n^2)$ operations. Consequently, we are interested in structure preserving properties of the double bracket.

Actually we do not predict that isospectral flows will be directly used to effectively find eigenvalues. However, we do hope that the study of such flows will lead to matrix factorizations which do so. For more on the relationship between isospectral flows and matrix factorizations see Chu [6], Watkins [27] and, Chu and Norris [11].

We are particularly interested in operators on matrices that preserve both spectrum and Toeplitz or near Toeplitz structure. There has been some study of such linear operators. Chu [10] obtained the following result. We use J to denote the $n \times n$ exchange matrix $[J_{ij}]$, where $J_{ij} = 1$ if $i + j = n + 1$ otherwise $J_{ij} = 0$, and I' to denote the matrix $\text{Diag}(-1, +1, -1, \dots, (-1)^{n-1})$.

Proposition 1.3. *Let Q be a real orthogonal matrix. Then the following conditions are equivalent:*

- (1) *For every real symmetric matrix A , if A is Toeplitz, then so is QAQ^t .*
- (2) *The matrix Q is one of the following: $\pm I, \pm J, \pm I', \pm I'J$.*

Driessel and So [18] obtained the following related result. For complex number λ , we use $D(\lambda)$ to denote $\text{Diag}(1, \lambda, \dots, \lambda^{n-1})$.

Proposition 1.4. *Let T be a nonzero linear map on the real space of Hermitian matrices. Then the following conditions are equivalent:*

(1) *For every Hermitian matrix A , the spectrum of A is the same as the spectrum of $T(A)$ and the displacement rank of A is the same as the displacement rank of $T(A)$.*

(2) *There is a complex number λ such that $|\lambda| = 1$ and either for all Hermitian A , $T(A) = D(\lambda)AD(\lambda)^*$ or for all Hermitian A , $T(A) = D(\lambda)A^tD(\lambda)^*$.*

These results show that there are very few linear operators preserving spectrum and Toeplitz structure. Consequently, we begin to study double operators that do so. Unfortunately, we shall find that there are also very few double bracket operators that do so.

2. Preserving Hermitian Tridiagonal Matrices

We shall say that $A \in \mathbb{C}^{n \times n}$ is a *Jacobi* matrix if A is Hermitian and tridiagonal. In this section we consider the following:

Problem A. Find all Hermitian matrices M with the “Jacobi preserving” property (J):

$$\text{if } A \text{ is Jacobi, then so is } [A, [A, M]].$$

Let $\mathcal{J}(n)$ denote the set of Hermitian matrices which satisfy the Jacobi preserving property; in symbols,

$$\mathcal{J}(n) := \{M \in \mathbb{C}^{n \times n} : M^* = M \text{ and } M \text{ satisfies } (J)\}.$$

For A , let $ad(A)$ denote the linear map $X \rightarrow [A, X]$ on $\mathbb{C}^{n \times n}$. In the theory of Lie algebras this map is called the *adjoint* map determined by A . Let $Jac(n)$ denote the set of Jacobi matrices with size n . Note that $Jac(n)$ is a real linear subspace of $\mathbb{C}^{n \times n}$ and hence that $ad(A)^{-1}(Jac(n)) = \{X \in \mathbb{C}^{n \times n} : [A, X] \in Jac(n)\}$ is also a real linear subspace. Note

further that a Hermitian matrix M satisfies property (J) iff for every $A \in Jac(n)$, $M \in (ad(A) \circ ad(A))^{-1}(Jac(n))$; in symbols,

$$\mathcal{J}(n) = \cap \{(ad(A) \circ ad(A))^{-1}(Jac(n)) : A \in Jac(n)\} \cap Herm(n),$$

where $Herm(n)$ is the set of all $n \times n$ Hermitian matrices. Thus we see that $\mathcal{J}(n)$ is the intersection of certain real subspaces of $\mathbb{C}^{n \times n}$ and, in particular, we see that $\mathcal{J}(n)$ is itself such a subspace. To solve the given problem we shall adopt the following strategy: We select $A \in Jac(n)$ for which it is easy to compute $ad(A)$. We then determine $(ad(A) \circ ad(A))^{-1}(Jac(n))$. In this way we reduce the possible forms for $M \in \mathcal{J}(n)$.

The simplest choice for A is a diagonal matrix. Let D be a diagonal matrix with real distinct diagonal entries d_1, d_2, \dots, d_n ; in symbols, $D = Diag(d_1, d_2, \dots, d_n)$. Then $[D, X](i, j) = (d_i - d_j)X(i, j)$ and $[D, [D, X]](i, j) = (d_i - d_j)^2 X(i, j)$. Thus, we see that if $[D, [D, M]] \in Jac(n)$, then M must be a Jacobi matrix. In other words, M must have the following form:

$$M = \begin{bmatrix} x_1 & y_1 & 0 & \cdots & 0 & 0 \\ \bar{y}_1 & x_2 & y_2 & \cdots & 0 & 0 \\ 0 & \bar{y}_2 & x_3 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & x_{n-1} & y_{n-1} \\ 0 & 0 & 0 & \cdots & \bar{y}_{n-1} & x_n \end{bmatrix}.$$

We can easily find nondiagonal test matrices which convince us that M must be a diagonal matrix. In fact, let E_{pq} denote the matrix with 1 in the (p, q) position and 0's elsewhere; in symbols, $E_{pq}(i, j) := \delta(p, i)\delta(q, j)$, where δ is the Kronecker delta function. Take $A := E_{11} + E_{23} + E_{32}$. Computing with 3×3 matrices shows what happens. We have (using * for "don't care" positions)

$$\begin{aligned}
[A, M] &= \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} \begin{bmatrix} x_1 & y_1 & 0 \\ \bar{y}_1 & x_2 & y_2 \\ 0 & \bar{y}_2 & x_3 \end{bmatrix} - \begin{bmatrix} x_1 & y_1 & 0 \\ \bar{y}_1 & x_2 & y_2 \\ 0 & \bar{y}_2 & x_3 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} \\
&= \begin{bmatrix} * & y_1 & 0 \\ 0 & * & * \\ \bar{y}_1 & * & * \end{bmatrix} - \begin{bmatrix} * & 0 & y_1 \\ \bar{y}_1 & * & * \\ 0 & * & * \end{bmatrix} = \begin{bmatrix} * & y_1 & -y_1 \\ -\bar{y}_1 & * & * \\ \bar{y}_1 & * & * \end{bmatrix}
\end{aligned}$$

and

$$\begin{aligned}
[A, [A, M]] &= \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} \begin{bmatrix} * & y_1 & -y_1 \\ -\bar{y}_1 & * & * \\ \bar{y}_1 & * & * \end{bmatrix} - \begin{bmatrix} * & y_1 & -y_1 \\ -\bar{y}_1 & * & * \\ \bar{y}_1 & * & * \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} \\
&= \begin{bmatrix} * & * & -y_1 \\ * & * & * \\ -\bar{y}_1 & * & * \end{bmatrix} - \begin{bmatrix} * & * & y_1 \\ * & * & * \\ \bar{y}_1 & * & * \end{bmatrix} = \begin{bmatrix} * & * & -2y_1 \\ * & * & * \\ -2\bar{y}_1 & * & * \end{bmatrix}.
\end{aligned}$$

We see that y_1 must be zero since $[A, [A, M]]$ is Jacobi. Now, take $A = E_{12} + E_{21} + E_{33}$, then we have

$$\begin{aligned}
[A, M] &= \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x_1 & y_1 & 0 \\ \bar{y}_1 & x_2 & y_2 \\ 0 & \bar{y}_2 & x_3 \end{bmatrix} - \begin{bmatrix} x_1 & y_1 & 0 \\ \bar{y}_1 & x_2 & y_2 \\ 0 & \bar{y}_2 & x_3 \end{bmatrix} \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} \\
&= \begin{bmatrix} * & * & y_2 \\ * & * & 0 \\ 0 & \bar{y}_2 & * \end{bmatrix} - \begin{bmatrix} * & * & 0 \\ * & * & y_2 \\ \bar{y}_2 & 0 & * \end{bmatrix} = \begin{bmatrix} * & * & y_2 \\ * & * & -y_2 \\ -\bar{y}_2 & \bar{y}_2 & * \end{bmatrix}
\end{aligned}$$

and

$$[A, [A, M]] = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} * & * & y_2 \\ * & * & -y_2 \\ -\bar{y}_2 & \bar{y}_2 & * \end{bmatrix} - \begin{bmatrix} * & * & y_2 \\ * & * & -y_2 \\ -\bar{y}_2 & \bar{y}_2 & * \end{bmatrix} \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$= \begin{bmatrix} * & * & -y_2 \\ * & * & * \\ -\bar{y}_2 & * & * \end{bmatrix} - \begin{bmatrix} * & * & y_2 \\ * & * & * \\ \bar{y}_2 & * & * \end{bmatrix} = \begin{bmatrix} * & * & -2y_2 \\ * & * & * \\ -2\bar{y}_2 & * & * \end{bmatrix}.$$

We see that y_2 must be zero since $[A, [A, M]]$ is Jacobi. Similarly, we see that the rest of the off-diagonal elements of M must be zero; in symbols, $y_1 = y_2 = \dots = y_{n-1} = 0$. In other words, $M = \text{Diag}(x_1, x_2, \dots, x_n)$.

Now, let Z be the matrix with ones on the first superdiagonal and zeros elsewhere; in symbols, $Z = E_{12} + E_{23} + \dots + E_{n-1, n}$. We take $A = Z + Z^t$. Note that for any matrix W ,

$$[A, W](i, j) = W(i-1, j) + W(i+1, j) - W(i, j-1) - W(i, j+1).$$

We have used the convention that $W(i, j) = 0$ if either i or j is not between 1 and n . Computing with 4×4 matrices we see what happens. We have

$$[A, M] = \begin{bmatrix} 0 & x_2 - x_1 & 0 & 0 \\ x_1 - x_2 & 0 & x_3 - x_2 & 0 \\ 0 & x_2 - x_3 & 0 & x_4 - x_3 \\ 0 & 0 & x_3 - x_4 & 0 \end{bmatrix}$$

and

$$[A, [A, M]] = \begin{bmatrix} * & * & x_3 - 2x_2 + x_1 & 0 \\ * & * & * & x_4 - 2x_3 + x_2 \\ x_3 - 2x_2 + x_1 & * & * & * \\ 0 & x_4 - 2x_3 + x_2 & * & * \end{bmatrix}.$$

Thus, in general, we see that the second differences $x_{k+1} - 2x_k + x_{k-1}$ for $k = 2, 3, \dots, n-1$ must be zero. It follows that there exist real numbers a and b such that $x_k = a + bk$ for $k = 1, 2, \dots, n$. In other words, $M = aI + bD$, where $D := \text{Diag}(1, 2, \dots, n)$. We have obtained the solution of Problem A.

Theorem 2.1. *Let $n \geq 3$ and let M be a Hermitian matrix. Then the double bracket operator determined by M has the Jacobi preserving property iff M is a real linear combination of I and $\text{Diag}(1, 2, \dots, n)$.*

Proof. It is easy to verify that I and D have the Jacobi preserving property. On the other hand, if M has the Jacobi preserving property, then M must have the given form according to the argument preceding the statement of the theorem.

3. Preserving Hermitian Toeplitz Matrices

Recall that a matrix $A \in \mathbb{C}^{n \times n}$ is "Toeplitz" if it has constant diagonals; in symbols, $A(i, j) = A(i+1, j+1)$ for all $i, j = 1, 2, \dots, n-1$. In this section, we consider the following:

Problem B. Find all Hermitian matrices M with the "Toeplitz preserving" property (T):

if A is Toeplitz and Hermitian, then so is $[A, [A, M]]$.

Let $\mathcal{T}(n)$ denote the set of Hermitian matrices which satisfy the Toeplitz preserving property. Let $\text{Toep}(n)$ denote the set of Hermitian Toeplitz matrices with size n . Note that $\text{Toep}(n)$ is a real linear subspace of $\mathbb{C}^{n \times n}$. Also, note that $M \in \mathcal{T}(n)$ iff for every $A \in \text{Toep}(n)$, $M \in (\text{ad}(A) \circ \text{ad}(A))^{-1}(\text{Toep}(n))$; in symbols,

$$\mathcal{T}(n) = \bigcap \{(\text{ad}(A) \circ \text{ad}(A))^{-1}(\text{Toep}(n)) : A \in \text{Toep}(n)\}.$$

As in the previous section, we will use test matrices from $\text{Toep}(n)$ to reduce the possible forms for M . We first turn to the real version of Problem B. We shall see that the solution of this problem will provide part of the solution of Problem B.

Lemma 3.1. *Let M be a real symmetric matrix, then the following properties for M are equivalent:*

- (1) *if A is real symmetric and Toeplitz, then so is $[A, [A, M]]$;*

(2) if A and B are real symmetric and Toeplitz, then so is $[A, [B, M]] + [B, [A, M]]$;

(3) if A_1, \dots, A_m form a basis for real symmetric matrices, then $[A_i, [A_j, M]] + [A_j, [A_i, M]]$ is real symmetric and Toeplitz.

Proof. (1) \Rightarrow (2) follows from the identity:

$$\begin{aligned} [A + B, [A + B, M]] &= [A, [A, M]] + [A, [B, M]] \\ &\quad + [B, [A, M]] + [B, [B, M]]. \end{aligned}$$

(2) \Rightarrow (3) is clear. For (3) \Rightarrow (1), consider a real symmetric matrix A . Since $\{A_i\}$ is a basis, $A = \sum_i a_i A_i$. Hence

$$[A, [A, M]] = \sum_{i,j} a_i a_j ([A_i, [A_j, M]] + [A_j, [A_i, M]])$$

is also real symmetric.

Again we let Z be the matrix with ones on the first superdiagonal and zeros elsewhere. Let $H_p = Z^p + (Z^p)^t$. Then the set $\{I\} \cup \{H_p : p = 1, 2, \dots, n-1\}$ is a basis for real symmetric matrices. Recall that the "exchange" matrix J and note that $JAJ = A$ for every real symmetric Toeplitz matrix A , hence $[A, J] = 0$.

Theorem 3.2. *Let M be a real symmetric matrix. Then M satisfies the real Toeplitz preserving property (TR):*

$$\text{if } A \text{ is real symmetric Toeplitz, then so is } [A, [A, M]]$$

if and only if M is a real linear combination of I and J .

Proof. (\Leftarrow) Let $M = aI + bJ$, where a and b are real numbers. Then $[A, M] = 0$ for every symmetric Toeplitz matrix A .

(\Rightarrow) **Claim 1.** If $[H_1, K]$ is Toeplitz and K is real skew-symmetric, then $K = 0$.

We shall outline the proof of this claim for 5×5 matrices. It will be clear that the argument applies to the general case. Assume $[H_1, K]$ is Toeplitz and that K is skew-symmetric. Then K has the following form:

$$K = \begin{bmatrix} 0 & x_1 & * & * & * \\ -x_1 & 0 & x_2 & * & * \\ * & -x_2 & 0 & x_3 & * \\ * & * & -x_3 & 0 & x_4 \\ * & * & * & -x_4 & 0 \end{bmatrix}.$$

We do not care about the $*$ entries until later. Since $[H_1, K]$ is Toeplitz and the trace of $[H_1, K]$ is 0 we see that the diagonal entries are zero. In other words we have $0 = -2x_1 = 2(x_1 - x_2) = 2(x_2 - x_3) = 2(x_3 - x_4) = 2x_4$. We conclude that $0 = x_1 = x_2 = x_3 = x_4$. Thus, K has the following form:

$$K = \begin{bmatrix} 0 & 0 & y_1 & * & * \\ 0 & 0 & 0 & y_2 & * \\ -y_1 & 0 & 0 & 0 & y_3 \\ * & -y_2 & 0 & 0 & 0 \\ * & * & -y_3 & 0 & 0 \end{bmatrix}.$$

We consider the entries of $[H_1, K]$ which lie on the first superdiagonal. We have $[H_1, K]_{i, i+1} = a$ for some real scalar a and we also have, by direct computation

i	$[H_1, K]_{i, i+1}$
1	$-y_1$
2	$y_1 - y_2$
3	$y_2 - y_3$
4	y_3

Adding we get $4a = \sum_{i=1}^4 [H_1, K]_{i, i+1} = 0$. We conclude that $a = 0$ and

so $y_1 = y_2 = y_3 = 0$. Clearly we can continue in this way showing that successive superdiagonals of K are zero.

Claim 2. Let M be a real symmetric matrix which satisfies the property (TR). Then $[H_p, M] = 0$ for $p = 1, 2, \dots, n-1$.

Since H_1 is real symmetric Toeplitz, $[H_1, [H_1, M]]$ is Toeplitz. Then, by Claim 1, $[H_1, M] = 0$ since $[H_1, M]$ is real skew-symmetric. Note that $\{I\} \cup \{H_p : p = 1, \dots, n-1\}$ forms a basis for real symmetric Toeplitz. Hence, by Lemma 3.1, $[H_1, [H_p, M]] + [H_p, [H_1, M]] = [H_1, [H_p, M]]$ is Toeplitz since M has (TR) property. Consequently, by Claim 1, $[H_p, M] = 0$ since $[H_p, M]$ is real skew-symmetric.

Claim 3. If M commutes with every real symmetric Toeplitz matrix, then M is a real linear combination of I and J .

Assume M commutes with every symmetric Toeplitz matrix. Decompose M as follows:

$$M = \begin{bmatrix} a & x^t & b \\ x & P & y \\ b & y^t & c \end{bmatrix},$$

where a, b and c are real scalars, x and y are real vectors with length $n-2$, and P is an $(n-2) \times (n-2)$ real symmetric matrix. By Claim 2, we have

$$\begin{aligned} 0 &= [H_{n-1}, M] \\ &= \begin{bmatrix} 0 & \dots & 1 \\ \vdots & & \vdots \\ 1 & \dots & 0 \end{bmatrix} \begin{bmatrix} a & x^t & b \\ x & P & y \\ b & y^t & c \end{bmatrix} - \begin{bmatrix} a & x^t & b \\ x & P & y \\ b & y^t & c \end{bmatrix} \begin{bmatrix} 0 & \dots & 1 \\ \vdots & & \vdots \\ 1 & \dots & 0 \end{bmatrix} \\ &= \begin{bmatrix} b & y^t & c \\ 0 & 0 & 0 \\ a & x^t & b \end{bmatrix} - \begin{bmatrix} b & 0 & a \\ y & 0 & x \\ c & 0 & b \end{bmatrix} = \begin{bmatrix} 0 & y^t & c-a \\ -y & 0 & -x \\ a-c & x^t & 0 \end{bmatrix}. \end{aligned}$$

We conclude that $a = c$ and $x = y = 0$. In other words, M has the following form

$$M = \begin{bmatrix} a & 0 & b \\ 0 & P & 0 \\ b & 0 & a \end{bmatrix}.$$

Let $N := M - aI - bJ$. Then N has the form

$$N = \begin{bmatrix} 0 & 0 & 0 \\ 0 & R & 0 \\ 0 & 0 & 0 \end{bmatrix},$$

where R is an $(n - 2) \times (n - 2)$ matrix. Note that N commutes with every symmetric Toeplitz matrix. We want to see that $R = 0$. Let A be an arbitrary symmetric Toeplitz matrix. Decompose A as follows:

$$A = \begin{bmatrix} p & u^t & q \\ u & T & Ju \\ q & (Ju)^t & p \end{bmatrix},$$

where p and q are scalars, u is a vector with length $n - 2$ and T is the $(n - 2) \times (n - 2)$ symmetric Toeplitz matrix determined by p and u . We consider $[A, N]$. We have

$$\begin{aligned} 0 &= \begin{bmatrix} p & u^t & q \\ u & T & Ju \\ q & (Ju)^t & p \end{bmatrix} \begin{bmatrix} 0 & 0 & 0 \\ 0 & R & 0 \\ 0 & 0 & 0 \end{bmatrix} - \begin{bmatrix} 0 & 0 & 0 \\ 0 & R & 0 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} p & u^t & q \\ u & T & Ju \\ q & (Ju)^t & p \end{bmatrix} \\ &= \begin{bmatrix} * & * & * \\ Ru & * & * \\ * & * & * \end{bmatrix}. \end{aligned}$$

From this we see that $Ru = 0$ for every vector u . Thus $R = 0$.

Lemma 3.3. *If the last three rows and columns of X are zero and $[H_1, [H_1, X]]$ is Toeplitz, then $X = 0$.*

Proof. We shall outline the proof of the claim for 6×6 matrices. It will be clear that the argument applies to the general case. Let

$$X = \begin{bmatrix} x_{11} & x_{12} & x_{13} & 0 & 0 & 0 \\ x_{21} & x_{22} & x_{23} & 0 & 0 & 0 \\ x_{31} & x_{32} & x_{33} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}.$$

We have

$$[H_1, X] = \begin{bmatrix} * & * & * & -x_{13} & 0 & 0 \\ * & * & * & -x_{23} & 0 & 0 \\ * & * & * & -x_{33} & 0 & 0 \\ x_{31} & x_{32} & x_{33} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

and

$$[H_1, [H_1, X]] = \begin{bmatrix} * & * & * & * & x_{13} & 0 \\ * & * & * & * & x_{23} & 0 \\ * & * & * & * & x_{33} & 0 \\ * & * & * & * & 0 & 0 \\ x_{31} & x_{32} & x_{33} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}.$$

Since this matrix is Toeplitz we conclude that $x_{13} = x_{23} = x_{33} = 0$ and $x_{31} = x_{32} = x_{33} = 0$. In other words, the last four rows and columns of X are zero. Clearly we can continue showing more rows and columns are zero. Finally, we get $X = 0$.

Theorem 3.4. *Let N be a real skew-symmetric matrix with dimension greater than 3. If N has the property (TS):*

if A is Hermitian Toeplitz, then so is $[A, [A, N]]$,

then $N = 0$.

Proof. Assume that N is a real skew-symmetric matrix which satisfies the property (TS). Again, we use appropriate test matrices to discover the form of N . We first use matrices H_p , where $p > n/2$. Note that such a matrix has the following block form:

$$H_p = \begin{bmatrix} 0 & 0 & I \\ 0 & 0 & 0 \\ I & 0 & 0 \end{bmatrix},$$

where $I = I_{n-p}$ is the $(n-p) \times (n-p)$ identity matrix. For any matrix X we partition as follows:

$$X = \begin{bmatrix} X_{11} & X_{12} & X_{13} \\ X_{21} & X_{22} & X_{23} \\ X_{31} & X_{32} & X_{33} \end{bmatrix}.$$

Then

$$[H_p, [H_p, X]] = \begin{bmatrix} 2(X_{11} - X_{33}) & X_{12} & 2(X_{13} - X_{31}) \\ X_{21} & 0 & X_{23} \\ 2(X_{31} - X_{13}) & X_{32} & 2(X_{33} - X_{11}) \end{bmatrix}.$$

In particular, when $p = n - 1$ we partition N as follows:

$$N = \begin{bmatrix} 0 & x^t & b \\ -x & N_1 & y \\ -b & -y^t & 0 \end{bmatrix},$$

where b is a scalar, x and y are vectors and N_1 is an $(n-2) \times (n-2)$ real skew-symmetric matrix. We get

$$[H_{n-1}, [H_{n-1}, N]] = \begin{bmatrix} 0 & x^t & 4b \\ -x & 0 & y \\ -4b & -y^t & 0 \end{bmatrix}.$$

This matrix must be Toeplitz. We conclude that there is a scalar a such that

$$x = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ a \end{bmatrix}, \quad y = \begin{bmatrix} a \\ 0 \\ \vdots \\ 0 \\ 0 \end{bmatrix}.$$

In other words, N has the following form:

$$N = \begin{bmatrix} 0 & 0 & \cdots & a & b \\ 0 & & & & a \\ \vdots & & N_1 & \vdots & \\ -a & & & & 0 \\ -b & -a & \cdots & 0 & 0 \end{bmatrix}.$$

When $p = n - 2$, we partition N as follows:

$$N = \begin{bmatrix} 0 & 0 & 0 & a & b \\ 0 & 0 & x^t & b' & a \\ 0 & -x & N_2 & y & 0 \\ -a & -b' & -y^t & 0 & 0 \\ -b & -a & 0 & 0 & 0 \end{bmatrix}.$$

We get

$$[H_{n-2}, [H_{n-2}, N]] = \begin{bmatrix} 0 & 0 & 0 & 4a & 2(b + b') \\ 0 & 0 & x^t & 2(b + b') & 4a \\ 0 & -x & 0 & y & 0 \\ -4a & -2(b + b') & -y^t & 0 & 0 \\ -2(b + b') & -4a & 0 & 0 & 0 \end{bmatrix}.$$

Since this matrix must be Toeplitz, we conclude $x = 0$, $y = 0$ and $b + b' = 0$. In other words, N has the following form:

$$N = \begin{bmatrix} 0 & 0 & 0 & \cdots & 0 & a & b \\ 0 & 0 & 0 & \cdots & 0 & -b & a \\ 0 & 0 & & & & 0 & 0 \\ \vdots & \vdots & & N_2 & & \vdots & \vdots \\ 0 & 0 & & & & 0 & 0 \\ -a & b & 0 & \cdots & 0 & 0 & 0 \\ -b & -a & 0 & \cdots & 0 & 0 & 0 \end{bmatrix}.$$

When $p = n - 3$, we partition N as follows:

$$N = \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & a & b \\ 0 & 0 & 0 & 0 & 0 & -b & a \\ 0 & 0 & 0 & x^t & c & 0 & 0 \\ 0 & 0 & -x & N_3 & y & 0 & 0 \\ 0 & 0 & -c & -y^t & 0 & 0 & 0 \\ -a & b & 0 & 0 & 0 & 0 & 0 \\ -b & -a & 0 & 0 & 0 & 0 & 0 \end{bmatrix}.$$

We get

$$[H_{n-3}, [H_{n-3}, N]] = \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 2a & 2(b+c) \\ 0 & 0 & 0 & 0 & 2a & -4b & 2a \\ 0 & 0 & 0 & x^t & 2(b+c) & 2a & 0 \\ 0 & 0 & 0 & 0 & y & 0 & 0 \\ 0 & -2a & -2(b+c) & -y^t & 0 & 0 & 0 \\ -2a & 4b & -2a & 0 & 0 & 0 & 0 \\ -2(b+c) & -2a & 0 & 0 & 0 & 0 & 0 \end{bmatrix}.$$

Since this matrix is Toeplitz, we conclude that $x = 0$, $y = 0$, $b + c = 0$, $a = 0$ and $b = 0$. In other words, N has the following form:

$$N = \begin{bmatrix} 0 & 0 & 0 & \cdots & 0 & 0 & 0 \\ 0 & 0 & 0 & \cdots & 0 & 0 & 0 \\ 0 & 0 & 0 & \cdots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & N_3 & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & 0 & 0 & 0 \\ 0 & 0 & 0 & \cdots & 0 & 0 & 0 \\ 0 & 0 & 0 & \cdots & 0 & 0 & 0 \end{bmatrix}.$$

Case 1. $n \geq 7$. Then $n - 3 > \frac{n}{2}$. By the above argument, the last three rows and columns are zero. Hence, by Lemma 3.3, $N = 0$.

Case 2. $n = 6$. Then $n - 2 > \frac{n}{2}$. As above using test matrices $H_{n-1} = H_5$ and $H_{n-2} = H_4$ we see that N has the following form:

$$N = \begin{bmatrix} 0 & 0 & 0 & 0 & a & b \\ 0 & 0 & 0 & 0 & -b & a \\ 0 & 0 & 0 & c & 0 & 0 \\ 0 & 0 & -c & 0 & 0 & 0 \\ -a & b & 0 & 0 & 0 & 0 \\ -b & -a & 0 & 0 & 0 & 0 \end{bmatrix}$$

and

$$[H_3, [H_3, N]] = \begin{bmatrix} 0 & 0 & 0 & 0 & 2a & 2(b+c) \\ 0 & 0 & 0 & 2a & -4b & 2a \\ 0 & 0 & 0 & 2(b+c) & 2a & 0 \\ 0 & -2a & -2(b+c) & 0 & 0 & 0 \\ -2a & 4b & -2a & 0 & 0 & 0 \\ -2(b+c) & -2a & 0 & 0 & 0 & 0 \end{bmatrix}.$$

Since this matrix must be Toeplitz we conclude that $b + c = 0$, $a = 0$ and $b = 0$. Thus $N = 0$.

Case 3. $n = 5$. Then $n - 2 > \frac{n}{2}$. As above using $H_{n-1} = H_4$ and $H_{n-2} = H_3$, we see that N has the following form:

$$N = \begin{bmatrix} 0 & 0 & 0 & a & b \\ 0 & 0 & 0 & -b & a \\ 0 & 0 & 0 & 0 & 0 \\ -a & b & 0 & 0 & 0 \\ -b & -a & 0 & 0 & 0 \end{bmatrix}.$$

Then

$$[H_1, [H_1, N]] = \begin{bmatrix} 0 & a & 3b & a & * \\ -a & 0 & -a & 6b & a \\ -3b & a & 0 & -a & 3b \\ -a & 6b & a & 0 & a \\ * & -a & -3b & -a & 0 \end{bmatrix}.$$

Since this matrix is Toeplitz we conclude $a = b = 0$ and hence $N = 0$.

Case 4. $n = 4$. Then $n - 1 > \frac{n}{2}$. As above using $H_{n-1} = H_3$, we see that N has the following form:

$$N = \begin{bmatrix} 0 & 0 & a & b \\ 0 & 0 & b' & a \\ -a & -b' & 0 & 0 \\ -b & -a & 0 & 0 \end{bmatrix}$$

and

$$[H_2, [H_2, N]] = \begin{bmatrix} 0 & 0 & 4a & 2(b + b') \\ 0 & 0 & 2(b + b') & 4a \\ -4a & -2(b + b') & 0 & 0 \\ -2(b + b') & -4a & 0 & 0 \end{bmatrix}.$$

Since this matrix must be Toeplitz we conclude that $b + b' = 0$. In other words, N has the following form:

$$N = \begin{bmatrix} 0 & 0 & a & b \\ 0 & 0 & -b & a \\ -a & b & 0 & 0 \\ -b & -a & 0 & 0 \end{bmatrix}.$$

Then we have

$$[H_1, [H_1, N]] = \begin{bmatrix} 0 & 4b & a & * \\ -4b & 0 & -8b & a \\ -a & 8b & 0 & 4b \\ * & -a & -4b & 0 \end{bmatrix}.$$

Since this matrix must be Toeplitz we conclude that $b = 0$. Now, we use

$$K_1 = \begin{bmatrix} 0 & 1 & 0 & 0 \\ -1 & 0 & 1 & 0 \\ 0 & -1 & 0 & 1 \\ 0 & 0 & -1 & 0 \end{bmatrix} \text{ as a test matrix. Then}$$

$$[iK_1, [iK_1, N]] = -a \begin{bmatrix} 2 & 0 & -1 & 0 \\ 0 & -2 & 0 & -1 \\ -1 & 0 & -2 & 0 \\ 0 & -1 & 0 & 2 \end{bmatrix}.$$

Since this matrix is Toeplitz we conclude that $a = 0$ and hence $N = 0$.

Case 5. $n = 3$. Then $n - 1 > \frac{n}{2}$. As above using $H_{n-1} = H_2$ we see that N has the following form:

$$N = \begin{bmatrix} 0 & a & b \\ -a & 0 & a \\ -b & -a & 0 \end{bmatrix}.$$

Let

$$H_1 = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}, \quad K_1 = \begin{bmatrix} 0 & 1 & 0 \\ -1 & 0 & 1 \\ 0 & -1 & 0 \end{bmatrix}, \quad \text{and} \quad K_2 = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ -1 & 0 & 0 \end{bmatrix}.$$

Then H_1 , iK_1 , iK_2 , $H_1 iK_1$ and $H_1 + iK_2$ are all Hermitian Toeplitz.

Hence

$$\begin{aligned} & [H_1 + iK_1, [H_1 + iK_1, N]] - [H_1, [H_1, N]] - [iK_1, [iK_1, N]] \\ &= [H_1, [iK_1, N]] + [iK_1, [H_1, N]] \\ &= ib[H_1, [K_1, K_2]] + ia[K_1, [H_1, K_1]] + ib[K_1, [H_1, H_2]] \\ &= ia[K_1, [H_1, K_1]] + ib([H_1, [K_1, K_2]] + [K_1, [H_1, K_2]]) \\ &= -2ia \begin{bmatrix} 0 & -1 & 0 \\ -1 & 0 & -1 \\ 0 & -1 & 0 \end{bmatrix} + ib \begin{bmatrix} -4 & 0 & 0 \\ 0 & 8 & 0 \\ 0 & 0 & -4 \end{bmatrix}. \end{aligned}$$

This matrix must be Toeplitz and that b must be zero. In other words, N has the form aK_1 . We also consider

$$\begin{aligned} & [H_1 + iK_2, [H_1 + iK_2, N]] - [H_1, [H_1, N]] - [iK_2, [iK_2, N]] \\ &= [H_1, [iK_2, N]] + [iK_2, [H_1, N]] \\ &= ia([H_1, [K_2, K_1]] - [K_2, [H_1, K_1]]) \\ &= ia \begin{bmatrix} 2 & 0 & 6 \\ 0 & -4 & 0 \\ 6 & 0 & 2 \end{bmatrix}. \end{aligned}$$

This matrix must be Toeplitz and that a must be zero. Hence $N = 0$.

Theorem 3.5. *Let M be an $n \times n$ Hermitian matrix, where $n \geq 3$. Then M satisfies the Toeplitz preserving property (T) iff M is a real scalar matrix.*

Proof. (\Leftarrow) Let $M = \alpha I$, where α is a real number. Then $[A, M] = 0$ for every matrix A . Of course, M has the Toeplitz preserving property.

(\Rightarrow) Let $\operatorname{Re} M$ and $\operatorname{Im} M$ be the real and imaginary parts of M ; in symbols, $(\operatorname{Re} M)(j, k) := \operatorname{Re}(M(j, k))$ and $(\operatorname{Im} M)(j, k) := \operatorname{Im}(M(j, k))$. In particular, $\operatorname{Re} M$ and $\operatorname{Im} M$ are $n \times n$ real matrices with $M = \operatorname{Re} M + i \operatorname{Im} M$. Since M is Hermitian, $\operatorname{Re}(M)$ is real symmetric and $\operatorname{Im}(M)$ is real skew-symmetric. Since M satisfies the property (T), $\operatorname{Re}(M)$ satisfies the property (TR) and $\operatorname{Im}(M)$ satisfies the property (TS). Now, by Theorem 3.2, $\operatorname{Re}(M) = aI + bJ$. And, by Theorem 3.3, $\operatorname{Im}(M) = 0$. It remains to show that $b = 0$. Suppose that $b \neq 0$, then $[iK_1, [iK_1, J]]$ is Toeplitz. Note that $Z^t = JZJ$ and

$$\begin{aligned} [K, J] &= [Z - JZJ, J] \\ &= ((Z - JZJ)J - J(Z - JZJ)) \\ &= 2(ZJ - JZ) = 2[Z, J]. \end{aligned}$$

Hence we have

$$\begin{aligned} [iK_1, [iK_1, J]] &= -[K_1, [K_1, J]] = -2[Z - JZJ, [Z, J]] \\ &= (-2)((Z - JZJ)(ZJ - JZ) - (ZJ - JZ)(Z - JZJ)) \\ &= (-2)((Z^2J - ZJZ - JZJZJ + JZJJZ) \\ &\quad - (ZJZ - ZJJZJ - JZ^2 + JZJZJ)) \\ &= (-2)(Z^2J - ZJZ - JZJZJ + JZ^2 - ZJZ \\ &\quad + Z^2J + JZ^2 - JZJZJ) \end{aligned}$$

$$\begin{aligned}
&= (-2)(2Z^2J - 2ZJZ - 2JZJZJ + 2JZ^2) \\
&= (-4)(Z^2J + JZ^2 - ZZ^tJ - JZZ^t).
\end{aligned}$$

This matrix is not Toeplitz; for example, when $n = 3$, we have

$$Z^2J = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{bmatrix} \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix},$$

$$JZ^2 = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix} \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix},$$

$$ZZ^tJ = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix},$$

$$JZZ^t = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix} \begin{bmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix},$$

and hence

$$Z^2J + JZ^2 - ZZ^tJ - JZZ^t = \begin{bmatrix} 1 & 0 & -1 \\ 0 & -2 & 0 \\ -1 & 0 & 1 \end{bmatrix}.$$

We conclude that $b = 0$. In other words, $Re(M)$ is a real scalar and so is M .

Acknowledgements

Driessel and So began this research during the academic year 1991-92 at the Institute for Mathematics and Its Applications (IMA) at the University of Minnesota. We thank the members of the IMA and the University of Minnesota Mathematics Department for their hospitality. In particular, we wish to thank Jack Conn, Charles A. McCarthy and Hans Weinberger for their suggestions and encouragement. In that academic year the IMA concentrated on linear algebra. We thank the organizers of that program. Driessel and Hentzel presented some of these results in the linear algebra seminar at Iowa State University during the 1994-95 academic year. We thank the participants in that seminar, especially Bryan Cain, for their interest and suggestions. We also want to say that we have been strongly influenced by the connection between homogeneous autonomous differential equations and nonassociative algebras; see, for example, Walcher [26]; we thank S. Walcher for his help in this area.

References

- [1] A. M. Bloch, Steepest descent, linear programming, and Hamiltonian flows, *Contemporary Math.* 114 (1990), 77-88.
- [2] A. M. Bloch, R. W. Brockett and T. S. Ratiu, A new formulation of the generalized Toda lattice equations and their fixed point analysis via the momentum map, *Bull. Amer. Math. Soc. (N.S.)* 23 (1990), 477-485.
- [3] R. W. Brockett, Dynamical systems that sort lists and solve linear programming problems, *Proceedings of the 27th IEEE Conference on Decision and Control* (1988), 799-804.
- [4] R. W. Brockett, Least squares matching problems, *Linear Alg. Appl.* 122/123/124 (1989), 761-777.
- [5] R. W. Brockett, Dynamical systems that sort lists and solve linear programming problems, *Linear Alg. Appl.* 146 (1991), 79-91.
- [6] M. T. Chu, The generalized Toda flow, the QR algorithm and the center manifold theory, *SIAM J. Alg. Disc. Math.* 5 (1984), 187-210.
- [7] M. T. Chu, A differential equation approach to the singular value decomposition of bidiagonal matrices, *Linear Alg. Appl.* 80 (1986), 71-79.
- [8] M. T. Chu and L. K. Norris, Isospectral flows and abstract matrix factorizations, *SIAM J. Numer. Anal.* 25 (1988), 1383-1391.

- [9] M. T. Chu and K. R. Driessel, The projected gradient method for least squares matrix approximations with spectral constraints, *SIAM J. Numer. Anal.* 27 (1990), 1050-1060.
- [10] M. T. Chu, The stability group of symmetric Toeplitz matrices, *Linear Alg. Appl.* 185 (1993), 119-123.
- [11] M. T. Chu, Scaled Toda-like flows, *Linear Alg. Appl.* 215 (1995), 261-274.
- [12] P. Deift, T. Nanda and C. Tomei, Ordinary differential equations for the symmetric eigenvalue problem, *SIAM J. Numer. Anal.* 20 (1983), 1-22.
- [13] P. Deift, J. Demmel, L.-C. Li and C. Tomei, The bidiagonal singular value decomposition and Hamiltonian mechanics, *SIAM J. Numer. Anal.* 28 (1991), 1463-1516.
- [14] K. R. Driessel, On isospectral gradient flows: solving eigenproblems using differential equations, in *Inverse Problems - Proceedings of the Conference held at the Mathematical Research Institute at Oberwolfach, Black Forest, May 18-24, 1986*, pp. 69-91, edited by John R. Cannon and Ulrich Hornung, Birkhäuser, 1986.
- [15] K. R. Driessel, On finding the eigenvalues and eigenvectors of a matrix by means of an isospectral gradient flow, Technical Report No. 541, Department of Mathematical Sciences, Clemson University, Clemson, South Carolina, 1987a.
- [16] K. R. Driessel, On finding the singular values and singular vectors of a matrix by means of an isosingular gradient flow, Technical Report 87-01, Department of Mathematics, Idaho State University, Pocatello, Idaho, 1987b.
- [17] K. R. Driessel, Some homogeneous spaces of matrices, Technical Report 91-01, Department of Mathematics, Idaho State University, Pocatello, Idaho, 1991.
- [18] K. R. Driessel and W. So, Linear operators on matrices: preserving spectrum and displacement structure, *SIAM J. Matrix Anal.* 16 (1995), 488-501.
- [19] G. Golub and C. F. Van Loan, *Matrix Computations*, Johns Hopkins Press, Baltimore, 1983.
- [20] T. Nanda, Differential equations and the *QR* algorithm, *SIAM J. Numer. Anal.* 22 (1983), 310-321.
- [21] B. N. Parlett, *The Symmetric Eigenvalue Problem*, Prentice-Hall, Englewood Cliffs, New Jersey, 1980.
- [22] W. W. Symes, Hamiltonian group actions and integrable systems, *Physica* 1D (1980), 339-374.
- [23] W. W. Symes, The *QR* algorithm and scattering for the finite non-periodic Toda lattice, *Physica* 4D (1982), 275-280.
- [24] J. A. Thorpe, *Elementary Topics in Differential Geometry*, Springer-Verlag, New York, 1979.
- [25] C. Tomei, The topology of isospectral manifolds of tridiagonal matrices, *Duke Math. J.* 51 (1984), 981-996.

- [26] S. Walcher, *Algebras and Differential Equations*, Hadronic Press, Palm Harbor, 1991.
- [27] D. S. Watkins, *Isospectral flows*, *SIAM Review* 26 (1984), 379-391.
- [28] H. S. Wilf, *An algorithm-inspired proof of the spectral theorem in E^n* , *Amer. Math. Monthly* 88 (1981), 49-50.



JP JOURNAL OF ALGEBRA, NUMBER THEORY AND APPLICATIONS

The journal is devoted to publishing original research papers and survey articles in algebra, number theory and their applications. A volume of the JPANTA will appear in three issues published annually in April, August and December.

Submission of Manuscripts: Manuscripts, in duplicate, may be submitted to the Managing Editor or to a member of the Editorial Board for consideration in the JPANTA at the address of the Publisher noted at the end [Papers prepared in LATEX/AMS-TEX or Microsoft Word (95/97/2000) may be submitted electronically at the e-mail address: kkazad@pphmj.com or editors@pphmj.com]. The paper must be typed only on one side in double spacing with a generous margin all round. An effort is made to publish a paper duly recommended by a referee within a period of six months.

A proof will be sent to the author submitting the paper, unless requested otherwise, without the original manuscript, for corrections.

Authors are requested to provide an abstract of not more than 200 words and latest Mathematics Subject Classification. Statements of Lemmas, Propositions and Theorems should be set in *italics* and references should be arranged in alphabetical order by the surname of the first author in the following style:

Books

- [1] R. A. Mollin, Quadratics, CRC Press, Boca Raton, New York, London, Tokyo, 1996.

Research papers

- [2] Masao Toyozumi, A note on the Dedekind eta function, JP Jour. Algebra, Number Theory & Appl. (to appear).
[3] Mingfen Wu and Xiang-Yun Xie, Prime radical theorems on ordered semigroups, JP Jour. Algebra, Number Theory & Appl. 1(1) (2001), 1-9.

Page Charges: Authors are requested to arrange page charges of their papers @ GB£ 18.00 (or equivalent in EUR) per page in Europe and @ US\$ 25.00 per page in the rest of world from their institutions/research grants. However, for authors in India this charge is Rs. 250.00 per page. Twenty-five reprints of a paper are provided to the authors ex-gratis. Additional sets of reprints may be ordered at the time of proof correction.

Copyright: It is assumed that the submitted manuscript has not been published and will not be simultaneously submitted or published elsewhere. By submitting a manuscript, the authors agree that the copyright for their articles is transferred to the publisher, if and when, the paper is accepted for publication.

The publisher cannot take the responsibility of any loss of manuscript. Therefore, authors are requested to maintain a copy at their end.

Subscription: The price for a copy of Volume 1 (2001) of the JPANTA is Rs. 2000.00 in India and GB£ 177.00 (or equivalent in EUR) in Europe and US\$ 250.00 in the rest of world plus postage.

Publisher: Pushpa Publishing House
Vijaya Niwas, 198, Mumfordganj
Allahabad 211002, India
Phone and Fax: 0091-532-642078
e-mail: arun@pphmj.com
<http://www.pphmj.com>

JP Journal of Algebra, Number Theory and Applications

ISSN: 0972-5555

Volume 1(2001)

Baba, Kiyoshi and Yoshida, Ken-ichi: Flatness and faithful flatness of a two-generator extension $R[\alpha, \beta]$ of a Noetherian domain R	...	47
Baba, Kiyoshi and Yoshida, Ken-Ichi: Equivalence classes on linear fractional transforms of anti-integral elements	...	195
Ballico, E.: The cancellation problem for projective embeddings	...	213
Beasley, LeRoy B.; Cheon, Gi-Sang; Park, Se-Won and Seol, Han-Guk: Nonnegative sign patterns which allow normality over Z	...	133
Cao, Chong-Guang: See Zhang, Xian	...	77
Cao, Chong-Guang: See Zhang, Xian	...	225
Cheon, Gi-Sang: See Beasley, LeRoy B.	...	133
Cho, Yong Uk: A generalization of strong regularity in near-rings	...	115
Driessel, Kenneth R.; Hentzel, Irvin R. and So, Wasin: On preserving structured matrices using double bracket operators: Tridiagonal and toeplitz matrices	...	87
Ewell, John A.: Representing numbers by sums of three squares	...	203
Hentzel, Irvin R.: See Driessel, Kenneth R.	...	87
Kanemitsu, Mitsuo and Matsuda, Ryûki: On seminormal semigroups and pseudo-valuation semigroups	...	25

Kim, Seokwoo: See Seo, Changho	...	167
Kobayashi, Shigeru and Yoshikawa, Ryugo: On the subresultant chain in view of the pseudodivision	...	143
Lin, D. H.: See Liu, Y. L.	...	19
Liu, Y. L. and Lin, D. H.: The structure of simple p -semisimple BCI-algebras	...	19
Matsuda, Ryûki: Note on closedness of a semigroup ring	...	11
Matsuda, Ryûki: See Kanemitsu, Mitsuo	...	25
Mizuno, Hirobumi and Sato, Iwao: L -functions of graph coverings	...	235
Mollin, R. A.: Quadratic Diophantine equations determined by continued fractions	...	57
Oda, Susumu and Yoshida, Ken-ichi: Extensions $R[\beta] \cap R[\beta^{-1}]$ and $R[\alpha] \cap R[\alpha^{-1}]$ are equal for any transformation $\beta = (c\alpha - d)/(a\alpha - b)$ of an anti- integral element α with $ad - bc \in R^\times$...	41
Park, Se-Won: See Beasley, LeRoy B.	...	133
Sato, Iwao: Decomposition formulas of zeta functions of graphs and digraphs	...	217
Sato, Iwao: See Mizuno, Hirobumi	...	235
Seo, Changho; Kim, Seokwoo and Sung, Yeoulouk: On the crypto-coprocessor technic based on the elliptic curve cryptographic over the field $GF(2^{163})$ for Internal-Type Memory	...	167
Seol, Han-Guk: See Beasley, LeRoy B.	...	133

	iii
So, Wasin: See Driessel, Kenneth R.	87
Spearman, Blair K. and Williams, Kenneth S.: Integers which are discriminants of bicyclic or cyclic quartic fields	179
Sung, Yeolouk: See Seo, Changho	167
Toyozumi, Masao: A note on the Dedekind eta function	125
Williams, Kenneth S.: See Spearman, Blair K.	179
Wu, Mingfen and Xie, Xiang-Yun: Prime radical theorems on ordered semigroups	1
Xie, Xiang-Yun: See Wu, Mingfen	1
Yoshida, Ken-ichi: See Oda, Susumu	41
Yoshida, Ken-ichi: See Baba, Kiyoshi	47
Yoshida, Ken-Ichi: See Baba, Kiyoshi	195
Yoshikawa, Ryugo: See Kobayashi, Shigeru	143
Zhang, Xian and Cao, Chong-Guang: Endomorphisms of two dimensional unitary group over complex number field	77
Zhang, Xian and Cao, Chong-Guang: Homomorphisms between matrix multiplicative semigroups	225